

On the width of collective excitations in chiral soliton models

Herbert WEIGEL^{*)}

Fachbereich Physik, Siegen University, D-57068 Siegen, Germany

In chiral soliton models for baryons the computation of hadronic decay widths of baryon resonances is a long standing problem. For the three flavor Skyrme model I present a solution to this problem that satisfies large- N_C consistency conditions. As an application I focus on the hadronic decay of the Θ and Θ^* pentaquarks.

§1. Statement of the problem

Hadronic decays of baryon resonances are commonly described by a Yukawa interaction of the generic structure

$$\mathcal{L}_{\text{int}} \sim g \bar{\psi}_{B'} \phi \psi_B, \quad (1.1)$$

where B' is the resonance that decays into baryon B and meson ϕ and g is a coupling constant. It is crucial that this interaction Lagrangian is *linear* in the meson field. If ϕ is a pseudoscalar meson this interaction yields the decay width $\Gamma(B' \rightarrow B\phi) \propto g^2 |\vec{p}_\phi|^3$, with $\vec{p}_\phi$ being the momentum of the outgoing meson.

The situation is significantly different in soliton models that are based on action functionals of only meson degrees of freedom, $\Gamma = \Gamma[\Phi]$. These action functionals contain classical (static) soliton solutions, Φ_{sol} , that are identified as baryons. The interaction of these baryons with mesons is described by the (small) meson fluctuations about the soliton: $\Phi = \Phi_{\text{sol}} + \phi$. By pure definition we have

$$\left. \frac{\delta \Gamma[\Phi]}{\delta \Phi} \right|_{\Phi=\Phi_{\text{sol}}} = 0. \quad (1.2)$$

Thus there is no term linear in ϕ to be associated with the Yukawa interaction, eq. (1.1). This puzzle has become famous as the Yukawa problem in soliton models. However, this does not mean that soliton models cannot describe resonance widths. On the contrary, these widths can be extracted from meson baryon scattering amplitudes, just as in experiment. In soliton models two-meson processes acquire contributions from the second order term

$$\Gamma^{(2)} = \frac{1}{2} \phi \cdot \left. \frac{\delta^2 \Gamma[\Phi]}{\delta^2 \Phi} \right|_{\Phi=\Phi_{\text{sol}}} \cdot \phi. \quad (1.3)$$

This expansion simultaneously represents an expansion in N_C , the number of color degrees of freedom: $\Gamma = \mathcal{O}(N_C)$ and $\Gamma^{(2)} = \mathcal{O}(N_C^0)$. Terms $\mathcal{O}(\phi^3)$ vanish in the limit $N_C \rightarrow \infty$. This implies that $\Gamma^{(2)}$ contains all large- N_C information about hadronic decays of resonances. We may reverse this statement to argue about any computation

^{*)} e-mail address: Weigel@physik.uni-siegen.de

of hadronic decay widths in soliton models: For $N_C \rightarrow \infty$ it *must* identically match the result obtained from $\Gamma^{(2)}$. Unfortunately, the most prominent baryon resonance, the Δ isobar, becomes degenerate with the nucleon as $N_C \rightarrow \infty$. It is stable in that limit and its decay is not subject to the above described litmus-test. The situation is more interesting in soliton models for flavor $SU(3)$. In the so-called rigid rotator approach (RRA), that generates baryon states as (flavor) rotational excitations of the soliton, exotic resonances emerge that dwell in the anti-decuplet representation of flavor $SU(3)$.¹⁾ The most discussed (and disputed) such state is the Θ^+ pentaquark with zero isospin and strangeness $S = +1$. In the limit $N_C \rightarrow \infty$ the (would-be) anti-decuplet states maintain a non-zero mass difference with respect to the nucleon. Therefore the properties of Θ^+ predicted from any model treatment must also be seen in the S -matrix for kaon-nucleon scattering as computed from $\Gamma^{(2)}$. This (seemingly alternative) quantization of strangeness degrees of freedom is called the bound state approach (BSA) because in the $S = -1$ sector the resulting equations of motion for ϕ yield a bound state. Its occupation serves to describe the ordinary hyperons, Λ , Σ , Σ^* , etc.. The above discussed litmus-test requires that the BSA and RRA give identical results for the Θ^+ properties as $N_C \rightarrow \infty$. This did not seem to be true and it was argued that the prediction of pentaquarks would be a mere artifact of the RRA.²⁾ Here we will show that this is a premature conclusion and that pentaquark states do indeed emerge in both approaches. Furthermore the comparison between the BSA and RRA provides an unambiguous computation of pentaquark widths: It differs substantially from previous approaches³⁾ that adopted transition operators for $\Theta^+ \rightarrow KN$ from the axial current.

This presentation is based on ref.⁴⁾ which the interested reader may want to consult for further details.

§2. The model

For simplicity we consider the Skyrme model⁵⁾ as a particular example for chiral soliton models. However, we stress that our qualitative results do indeed generalize to *all* chiral soliton models because these results solely reflect the treatment of the model degrees of freedom.

Chiral soliton models are functionals of the chiral field, U , the non-linear realization of the pseudoscalar mesons^{*)}, ϕ_a

$$U(\vec{x}, t) = \exp \left[\frac{i}{f_\pi} \phi_a(\vec{x}, t) \lambda_a \right], \quad (2.1)$$

with λ_a being the Gell-Mann matrices of $SU(3)$. For a convenient presentation of the model we split the action into three pieces

$$\Gamma = \Gamma_{SK} + \Gamma_{WZ} + \Gamma_{SB}. \quad (2.2)$$

*) A remark on notation: In what follows we adopt the convention that repeated indices are summed over in the range $a, b, c, \dots = 1, \dots, 8$, $\alpha, \beta, \gamma, \dots = 4, \dots, 7$ and $i, j, k, \dots = 1, 2, 3$.

The first term represents the Skyrme model action

$$\Gamma_{SK} = \int d^4x \operatorname{tr} \left\{ \frac{f_\pi^2}{4} [\partial_\mu U \partial^\mu U^\dagger] + \frac{1}{32\epsilon^2} [U^\dagger \partial_\mu U, U^\dagger \partial_\nu U]^2 \right\}. \quad (2.3)$$

Here $f_\pi = 93\text{MeV}$ is the pion decay constant and ϵ is the dimensionless Skyrme parameter. In principle this is a free model parameter. The two-flavor version of the Skyrme model suggests to put $\epsilon = 4.25$ from reproducing the Δ -nucleon mass difference^{*)}. The QCD anomaly is incorporated via the Wess–Zumino action⁷⁾

$$\Gamma_{WZ} = -\frac{iN_C}{240\pi^2} \int d^5x \epsilon^{\mu\nu\rho\sigma\tau} \operatorname{tr} [\alpha_\mu \alpha_\nu \alpha_\rho \alpha_\sigma \alpha_\tau], \quad (2.4)$$

with $\alpha_\mu = U^\dagger \partial_\mu U$. Note that Γ_{WZ} vanishes in the two-flavor version of the model.

The flavor symmetry breaking terms are contained in Γ_{SB}

$$\Gamma_{SB} = \frac{f_\pi^2}{4} \int d^4x \operatorname{tr} [\mathcal{M} (U + U^\dagger - 2)], \quad \mathcal{M} = \operatorname{diag} (m_\pi^2, m_\pi^2, 2m_K^2 - m_\pi^2). \quad (2.5)$$

We do not include terms that distinguish between pion and kaon decay constants even though they differ by about 20% empirically. This omission is a matter of convenience and leads to an underestimation of symmetry breaking effects⁸⁾ which approximately can be accounted for by rescaling the kaon mass $m_K \rightarrow m_K f_K / f_\pi$.

The action, eq. (2.2) allows for a topologically non-trivial classical solution, the famous hedgehog soliton

$$\Phi_{\text{sol}} \sim U_0(\vec{x}) = \exp [i\vec{\lambda} \cdot \hat{x} F(r)], \quad r = |\vec{x}| \quad (2.6)$$

embedded in the isospin subspace of flavor $SU(3)$. The chiral angle, $F(r)$ solves the classical equation of motion subject to the boundary condition $F(0) - F(\infty) = \pi$ ensuring unit winding (baryon) number. The soliton can be constructed as a function of the dimensionless variable $\epsilon f_\pi r$ and is thus not subject to N_C scaling.

In the RRA baryon states are generated by canonically quantizing collective coordinates $A \in SU(3)$ that describe the (spin) flavor orientation of the soliton, $A(t)U_0(\vec{x})A^\dagger(t)$. The resultant eigenstates may be classified according to $SU(3)$ multiplets; see ref.⁹⁾ for a review.

§3. Large N_C P -wave channel phase shifts with strangeness

As motivated after eq. (1.3) we introduce fluctuations $\phi \sim \eta_\alpha(\vec{x}, t)$

$$U(\vec{x}, t) = \sqrt{U_0(\vec{x})} \exp \left[\frac{i}{f_\pi} \lambda_\alpha \eta_\alpha(\vec{x}, t) \right] \sqrt{U_0(\vec{x})}, \quad (3.1)$$

for the kaon fields.¹⁰⁾ Expanding the action in powers of these fluctuations is an expansion in η_α/f_π and thus a systematic series in $1/\sqrt{N_C}$. The term quadratic

^{*)} To ensure that the (perturbative) n -point functions scale as $N_C^{1-n/26}$ we substitute $f_\pi = 93\text{MeV} \sqrt{N_C/3}$ and $\epsilon = 4.25 \sqrt{3/N_C}$ in the study of the N_C dependence.

in η_α describes meson scattering off a potential generated by the classical soliton, eq. (2.6). The P -wave mode is characterized by a single radial function

$$\begin{pmatrix} \eta_4 + i\eta_5 \\ \eta_6 + i\eta_7 \end{pmatrix}_P(\vec{x}, t) = \int_{-\infty}^{\infty} d\omega e^{i\omega t} \eta_\omega(r) \hat{x} \cdot \vec{\tau} \chi(\omega). \quad (3.2)$$

In future we will omit the subscript that indicates the Fourier frequency. Upon quantization the components of the two-component iso-spinor $\chi(\omega)$ are elevated to creation- and annihilation operators. It is straightforward to deduce the Schrödinger type equation

$$h^2 \eta(r) + \omega [2\lambda(r) - \omega M_K(r)] \eta(r) = 0 \quad \text{with} \quad h^2 = -\frac{d^2}{dr^2} - \frac{2}{r} \frac{d}{dr} + V_{\text{eff}}(r). \quad (3.3)$$

The radial functions arise from the chiral angle $F(r)$ and may be readily taken from the literature.¹⁰⁾ The equation of motion (3.3) is not invariant under particle conjugation $\omega \leftrightarrow -\omega$, and thus different for kaons ($\omega > 0$) and anti-kaons ($\omega < 0$). This difference stems from the Wess-Zumino term. Equation (3.3) has a bound state solution at $\omega = -\omega_A$, that gives the mass difference between the Λ -hyperon and the nucleon in the large- N_C limit. As this energy eigenvalue is negative it corresponds to a kaon, *i.e.* it carries strangeness $S = -1$. In the symmetric case ($m_K = m_\pi$) the bound state is simply the zero mode of $SU(3)$ flavor symmetry. The WZ-term moves the potential bound state with $S = +1$ to the positive continuum and we expect a resonance structure in that channel. The corresponding phase shift is shown in the left panel of figure 1. No clear resonance structure is visible; the phase shifts hardly reach $\pi/2$. The absence of such a resonance has previously lead to the premature criticism that there would not exist a bound pentaquark in the large- N_C limit.²⁾

§4. Constraint fluctuations

To study the coupling between the fluctuations and the collective excitations we generalize eq. (3.1) to

$$U(\vec{x}, t) = A(t) \sqrt{U_0(\vec{x})} \exp \left[\frac{i}{f_\pi} \lambda_\alpha \tilde{\eta}_\alpha(\vec{x}, t) \right] \sqrt{U_0(\vec{x})} A^\dagger(t). \quad (4.1)$$




Fig. 1. Large N_C P -wave phase shifts with strangeness $S = +1$ as function of the kaon momentum. Left panel: unconstrained; right panel: constrained to be orthogonal to the collective rotation.

These fluctuations dwell in the intrinsic system as they rotate along with the soliton. The kaon P -wave is subject to the modified integro-differential equation

$$h^2 \tilde{\eta}(r) + \omega [2\lambda(r) - \omega M_K(r)] \tilde{\eta}(r) = -z(r) \left[\int_0^\infty r'^2 dr' z(r') 2\lambda(r') \tilde{\eta}(r') \right] \\ \times \left[2\lambda(r) - (\omega + \omega_0) M_K(r) - \omega_0 \left(\frac{X_\Theta^2}{\omega_\Theta - \omega} + \frac{X_\Lambda^2}{\omega} \right) (2\lambda(r) - \omega_0 M_K(r)) \right], \quad (4.2)$$

for the flavor symmetric case^{*)}. The radial function $\tilde{\eta}(r)$ is defined according to eq. (3.2) and $z(r) = \sqrt{4\pi} \frac{f_\pi}{\sqrt{\Theta_K}} \sin \frac{F(r)}{2}$ is the collective mode wave-function normalized with respect to the moment of inertia for flavor rotations into strangeness direction, $\Theta_K = f_\pi^2 \int d^3r M_K(r) \sin^2 \frac{F(r)}{2} = \mathcal{O}(N_C)$. The non-local terms without $X_{\Lambda, \Theta}$ reflect the constraint $\int dr r^2 z(r) M_K(r) \tilde{\eta}(r) = 0$ which avoids double counting of rotational modes in strangeness direction. The interesting coupling is contained in the interaction Hamiltonian

$$H_{\text{int}} = \frac{2}{\sqrt{4\pi\Theta_K}} d_{i\alpha\beta} D_{\gamma\alpha} R_\beta \int d^3r z(r) [2\lambda(r) - \omega_0 M_K(r)] \hat{x}_i \tilde{\xi}_\gamma(\vec{x}, t), \quad (4.3)$$

where $\tilde{\xi}_a = D_{ab} \tilde{\eta}_b$ are the fluctuations in the laboratory frame, that we actually detect in KN scattering. The collective coordinates are parameterized via the adjoint representation $D_{ab}(A) = \frac{1}{2} \text{tr} [\lambda_a A \lambda_b A^\dagger]$ and the $SU(3)$ generators R_a . Integrating out the collective degrees of freedom by means of standard perturbation theory induces the separable potential

$$\frac{|\langle \Theta | H_{\text{int}} | (KN)_{I=0} \rangle|^2}{\omega_\Theta - \omega} + \frac{|\langle \Lambda | H_{\text{int}} | (KN)_{I=0} \rangle|^2}{\omega_\Lambda + \omega}. \quad (4.4)$$

These matrix elements concern the T -matrix elements in the laboratory frame. Since the laboratory and intrinsic T -matrix elements are identical for the Θ^+ channel,¹¹⁾ we may add the exchange potential, eq. (4.4) in the intrinsic frame. We define matrix elements of collective coordinate operators

$$\langle \Theta^+ | d_{3\alpha\beta} D_{+\alpha} R_\beta | n \rangle =: X_\Theta \sqrt{\frac{N_C}{32}} \quad \text{and} \quad \langle \Lambda | d_{3\alpha\beta} D_{-\alpha} R_\beta | p \rangle =: X_\Lambda \sqrt{\frac{N_C}{32}}, \quad (4.5)$$

to end up with eq. (4.2). The first factor in the coefficient $\omega_0 = 2 \left(\frac{2}{\sqrt{\Theta_K}} \sqrt{\frac{N_C}{32}} \right)^2 = \frac{N_C}{4\Theta_K}$ arises in the equation of motion because the potential, eq. (4.4) is quadratic in the fluctuations. The remaining (squared) factors stem from the definitions of $X_{\Theta, \Lambda}$ and the constant of proportionality in H_{int} . The $X_{\Theta, \Lambda}$ must be computed with the methods provided in ref.¹²⁾ but generalized to arbitrary (odd) N_C .⁴⁾ For $N_C \rightarrow \infty$ we have $X_\Theta \rightarrow 1$ and $X_\Lambda \rightarrow 0$. From the orthogonality conditions of the equation of motion (3.3) we straightforwardly verify that

$$\tilde{\eta}(r) = \eta(r) - az(r) \quad \text{with} \quad a = \int_0^\infty dr r^2 z(r) M_K(r) \eta(r). \quad (4.6)$$

^{*)} The more complicated case $m_K \neq m_\pi$ is at length discussed in ref.⁴⁾

solves eq. (4.2) for large N_C . This is essential because, as $z(r)$ is localized in space, η and $\tilde{\eta}$ have identical phase shifts! Hence the litmus-test discussed in the introduction is indeed satisfied. The physics behind $\tilde{\eta}$ is best understood when introducing reduced wave-functions $\bar{\eta}(r)$ that solve eq. (4.2) modified with $X_\Theta \equiv X_\Lambda \equiv 0$. *i. e.* the collective excitations are decoupled. These wave-functions are still orthogonal to the collective modes and actually lead to the background phase shift shown in figure 1. Having obtained the $\bar{\eta}(r)$ we may again switch on the exchange contributions, eq. (4.4). The additional separable potential is treated by standard R-matrix techniques and augments the phase shift by

$$\tan(\delta_R(k)) = \frac{\Gamma(\omega_k)/2}{\omega_\Theta - \omega_k + \Delta(\omega_k)}. \quad (4.7)$$

Here $\omega_\Theta = \frac{N_C+3}{4\Theta_K}$ is the RRA result for the excitation energy of Θ . This phase shift exhibits the canonical resonance structure with the width and the pole shift

$$\Gamma(\omega_k) = 2k\omega_0 X_\Theta^2 \left| \int_0^\infty r^2 dr z(r) 2\lambda(r) \bar{\eta}_{\omega_k}(r) \right|^2, \quad (4.8)$$

$$\Delta(\omega_k) = \frac{1}{2\pi\omega_k} \mathcal{P} \int_0^\infty q dq \left[\frac{\Gamma(\omega_q)}{\omega_k - \omega_q} + \frac{\Gamma(-\omega_q)}{\omega_k + \omega_q} \right], \quad (4.9)$$

respectively. We have numerically verified that in the large- N_C limit with $X_\Theta^2 = 1$, the phase shift from eq. (4.7) is identical to what is labeled resonance phase shift in figure 1, that we calculated as the difference between the total (η) and background ($\bar{\eta}$) phase shifts. For finite N_C we have $X_\Theta \neq 1$ and $X_\Lambda \neq 0$ so the R-matrix formalism becomes two-dimensional (Λ and Θ^+ exchange). Contrary to earlier criticisms²⁾ the large N_C pentaquark phase shift indeed resonates!

§5. Results

In figure 2 we show the resonance phase shift computed from eq. (4.7) for various values of N_C . First we observe that the resonance position quickly moves towards larger energies as N_C decreases. This is mainly due to the strong N_C dependence of ω_Θ : For $N_C = 3$ it is twice as large as in the limit $N_C \rightarrow \infty$. The pole shift Δ is actually quite small (some ten MeV) so that ω_Θ is indeed a reliable estimate of the resonance energy. Second, the resonance becomes shaper as N_C decreases. To major parts this is caused by the reduction of X_Θ .

We now turn to more quantitative results for which we also include flavor symmetry breaking effects. Then the resonance position changes to

$$\omega_\Theta = \frac{1}{2} \left[\sqrt{\omega_0^2 + \frac{3\Gamma}{2\Theta_K}} + \omega_0 \right] + \mathcal{O}\left(\frac{1}{N_C}\right). \quad (5.1)$$

where $\Gamma = \mathcal{O}(N_C)$ is a functional of the soliton that is proportional to the meson mass difference, $m_K^2 - m_\pi^2$. The $\mathcal{O}(1/N_C)$ piece is sizable for $N_C = 3$ and we compute it in the scenario of ref.¹²⁾ We then find $\omega_\Theta \approx 700\text{MeV}$; taking model dependencies




Fig. 2. The resonance phase shift as a function of N_C for $m_K = m_\pi$.

into account we expect the pentaquark to be about 600...900MeV heavier than the nucleon.

For the width calculations there are two principle differences. First, the interaction Hamiltonian acquires an additional term

$$H_{\text{int}}^{\text{sb}} = (m_K^2 - m_\pi^2) d_{i\alpha\beta} D_{\gamma\alpha} D_{8\beta} \int d^3r z(r) \gamma(r) \tilde{\xi}_\gamma(\vec{x}, t) \hat{x}_i, \quad (5.2)$$

The radial function $\gamma(r)$ is again given in terms of the chiral angle.⁴⁾ Second, the X_A does not vanish as $N_C \rightarrow \infty$ and the R -matrix formalism is always two dimensional. Nevertheless, the large- N_C solution is always of the form (4.6) and the BSA phase shift is recovered. The width function turns to

$$\Gamma(\omega_k) = 2k\omega_0 \left| \int_0^\infty r^2 dr z(r) \left[2X_\Theta \lambda(r) + \frac{Y_\Theta}{\omega_0} (m_K^2 - m_\pi^2) \right] \bar{\eta}_{\omega_k}(r) \right|^2, \quad (5.3)$$

where X_Θ and $Y_\Theta = \sqrt{8N_C/3} \langle \Theta^+ | d_{3\alpha\beta} D_{+\alpha} D_{8\beta} | n \rangle$ are to be computed in the RRA approach with full inclusion of flavor symmetry breaking effects.

This width function is shown (for $N_C = 3$) in figure 3 for Θ and its isovector partner Θ^* . The latter merely requires the appropriate modification of the matrix elements in eq. (4.5). Most importantly, the k^3 behavior of the width function, as suggested by the model, eq. (1.1) is reproduced only right above threshold, afterwards it levels off. Second, and somewhat surprising, the width of the non-ground state pentaquark is smaller than that of the lowest lying pentaquark. Our particular model yields $\Gamma_\Theta \approx 40\text{MeV}$ and $\Gamma_{\Theta^*} \approx 20\text{MeV}$. We note that there are certainly model ambiguities in these results.




Fig. 3. Model prediction for the width, $\Gamma(\omega)$ of Θ^+ (left) and Θ^{*+} (right) for $N_C = 3$ as function of the momentum $k = \sqrt{\omega^2 - m_K^2}$ for three values of the kaon mass. Note the unequal scales.

§6. Conclusions

We have discussed the chiral soliton model approach to KN scattering in the $S = +1$ channel which contains the potential¹⁴⁾ Θ^+ pentaquark, a state predicted as a flavor rotational excitation of the soliton. Though the exactly known large N_C phase shift suggests otherwise, the Θ emerges as a genuine resonance. Our central result is the width function for $\Theta \rightarrow KN$. In the flavor symmetric case it contains only a *single* collective coordinate operator and is thus very different from estimates that extract an effective Yukawa coupling from the axial current matrix element.³⁾ Since our approach matches the exact large N_C result, we must conclude that those axial current scenarios are erroneous¹³⁾ and that the cancellation among contributions to this matrix element is an invalid argument for a small pentaquark width.

Acknowledgments

This key note is based on a collaboration with H. Walliser, whose contribution is highly appreciated. I am very grateful to the organizers of this workshop for the invitation to contribute to the proceedings despite I was unable to participate.

References

- 1) L. C. Biedenharn and Y. Dothan, (1984), Print-84-1039 (DUKE);
M. Chemtob, Nucl. Phys. **B256** (1985) 600;
M. Praszalowicz, "SU(3) Skyrmion", in *Skyrmions and Anomalies*, (1987) p. 112;
H. Walliser, Nucl. Phys. **A548** (1992) 649;
D. Diakonov, V. Petrov, M. Polyakov, Z. Phys. **A359** (1997) 305;
H. Weigel, Eur. Phys. J. A **2** (1998) 391;
H. Walliser and V. B. Kopeliovich, J. Exp. Theor. Phys. **97**, 433 (2003);
J. R. Ellis, M. Karliner, M. Praszalowicz, JHEP **05** (2004) 002.
- 2) N. Itzhaki *et al.*, Nucl. Phys. **B684** (2004) 264;
T. D. Cohen, Phys. Lett. **B581** (2004) 175; Phys. Rev. **D70** (2004) 014011;
A. Cherman *et al.*, Phys. Rev. **D72** (2005) 094015.
- 3) Diakonov *et al.* and Ellis *et al.* in ref.¹⁾;
M. Praszalowicz, Acta Phys. Polon. **B35** (2004) 1625;
M. Praszalowicz and K. Goeke, Acta Phys. Polon. **B36** (2005) 2255;
G. S. Yang, H. C. Kim, K. Goeke, arXiv:hep-ph/0701168.
- 4) H. Walliser, H. Weigel, Eur. Phys. J. A **26** (2005) 361;
H. Weigel, arXiv:hep-ph/0510111; Eur. Phys. J. **A31** (2007) 495.
- 5) T. H. R. Skyrme, Proc. Roy. Soc. Lond. **A260** (1961) 127; Nucl. Phys. **31** (1962) 556.
- 6) G. 't Hooft, Nucl. Phys. **B72** (1974) 461;
E. Witten, Nucl. Phys. **B160** (1979) 57.
- 7) E. Witten, Nucl. Phys. **B223** (1983) 422; 433.
- 8) H. Weigel *et al.*, Phys. Rev. **D42** (1990) 3177.
- 9) H. Weigel, Int. J. Mod. Phys. **A11** (1996) 2419.
- 10) C. Callan and I. Klebanov, Nucl. Phys. **B262** (1985) 365,
C. Callan, K. Hornbostel, I. Klebanov, Phys. Lett. **B202** (1988) 296,
I. Klebanov, "Strangeness in the Standard Skyrme Model", in *Hadrons and Hadronic Matter* (1989) p. 223;
C. L. Schat, N. N. Scoccola, C. Gobbi, Nucl. Phys. **A585** (1995) 627.
- 11) A. Hayashi *et al.*, Phys. Lett. **B147** (1984) 5.
- 12) H. Yabu and K. Ando, Nucl. Phys. **B301** (1988) 601.
- 13) H. Weigel, arXiv:hep-ph/0703072.
- 14) The current experimental situation is summarized in: K. Hicks, arXiv:hep-ph/0703004.